

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
1 April 2004 (01.04.2004)

PCT

(10) International Publication Number
WO 2004/026831 A1

(51) International Patent Classification⁷: **C07D 209/30**,
A61K 31/404, A61P 43/00, C07D 401/04

(21) International Application Number:
PCT/EP2003/009969

(22) International Filing Date:
8 September 2003 (08.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/411,480 17 September 2002 (17.09.2002) US

(81) Designated States (*national*): AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: **F. HOFFMANN-LA ROCHE AG** [CH/CH];
124 Grenzacherstrasse, CH-4070 Basle (CH).

(72) Inventors: **MADERA, Ann, Marie**; 7892 Gate Way,
Dublin, CA 94568 (US). **WEIKERT, Robert, James**;
1027 Doeg Road, Boulder Creek, CA 95006 (US).


(74) Agent: **WAECHTER, Dieter**; 124 Grenzacherstrasse,
CH-4070 Basle (CH).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 2,4-SUBSTITUTED INDOLES AND THEIR USE AS 5-HT₆ MODULATORS



(I)

(57) Abstract: The present invention provides a compound of the formula (I): a pharmaceutically acceptable salt or a prodrug thereof, where R¹, R², R³, R⁴, p and n are those defined here. The present invention also provides compositions comprising, methods for using, and methods for preparing Compound of Formula (I).

2,4-SUBSTITUTED INDOLES AND THEIR USE AS 5-HT₆ MODULATORS


This invention relates to 2-substituted indoles, and associated compositions, methods for use as therapeutic agents, and methods of preparation thereof.

The actions of the neurotransmitter 5-hydroxytryptamine (5-HT) as a major modulatory neurotransmitter in the brain, are mediated through a number of receptor families termed 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆, and 5-HT₇. Based on a high level of 5-HT₆ receptor mRNA in the brain, it has been stated that the 5-HT₆ receptor may play a role in the pathology and treatment of central nerve system disorders. In particular, 5-HT₆ selective ligands have been identified as potentially useful in the treatment of certain CNS disorders such as Parkinson's disease, Huntington's disease, anxiety, depression, manic depression, psychoses, epilepsy, obsessive compulsive disorders, migraine, Alzheimer's disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia and bulimia, panic attacks, attention deficit hyperactivity disorder (ADHD), attention deficit disorder (ADD), withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Such compounds are also expected to be of use in the treatment of certain gastrointestinal (GI) disorders such as functional bowel disorder. See for example, B.L. Roth et al., *J. Pharmacol. Exp. Ther.*, 1994, 268, pages 1403-14120, D. R. Sibley et al., *Mol. Pharmacol.*, 1993, 43, 320-327, A.J. Sleight et al., *Neurotransmission*, 1995, 11, 1-5, and A. J. Sleight et al., *Serotonin ID Research Alert*, 1997, 2(3), 115-8. 5-HT₆ antagonists have also been identified as potentially useful compounds for treatment of obesity. See for example, Bentley et al., *Br. J. Pharmac.* 1999, *Suppl* 126; Bentley et al., *J. Psychopharmacol.* 1997, *Suppl* A64: 255; Wooley et al., *Neuropharmacology* 2001, 41: 210-129; and WO 02/098878.

While some 5-HT₆ modulators have been disclosed, there continues to be a need for compounds that are useful for modulating 5-HT₆.

- 2 -

One object of the present invention is (i) A compound of the formula:



or a pharmaceutically acceptable salt thereof,

5 wherein

n is 0, 1 or 2;

p is 1 or 2;

R¹ is optionally substituted aryl or optionally substituted heteroaryl;

R² is a optionally substituted heterocyclyl;

- 10 R³ is hydrogen, alkyl, or -C(=O)-R⁵, where R⁵ is alkyl, alkoxy, aryl, or aryloxy; and each R⁴ is independently hydrogen, hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino),
- 15 alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy.

Further objects of the present invention are:

(ii) The compound of (i),

wherein

n is 2;

- 20 R¹ is optionally substituted aryl or optionally substituted heteroaryl;

R² is a optionally substituted heterocyclyl;

R³ is hydrogen, alkyl;

and

R⁴ is hydrogen.


- 25 (iii) The compound of (ii), wherein R² is optionally substituted piperazin-1-yl or optionally substituted piperidin-4-yl.

(iv) The compound of (iii), wherein R² is piperazin-1-yl, 4-methylpiperazin-1-yl, 3,5-dimethylpiperazin-1-yl, N-methyl piperidin-4-yl or piperidin-4-yl.

- 3 -

- (v) The compound of (iv), wherein R^2 is piperazin-1-yl, 4-methylpiperazin-1-yl or piperidin-4-yl.
- (vi) The compound of (ii), wherein R^1 is optionally substituted phenyl or optionally substituted thienyl.
- 5 (vii) The compound (vi), wherein R^1 is thien-2-yl or phenyl which is optionally substituted with alkyl, halo, or haloalkyl.
- (viii) The compound of (vii), wherein R^1 is phenyl, 2,3-dichlorophenyl, 2-fluorophenyl, 2-methylphenyl, 2-trifluoromethylphenyl, 3-bromophenyl, or thien-2-yl.
- (ix) The compound of (ii), wherein R^3 is hydrogen or methyl.
- 10 (x) The compounds (ii), which are:
- 2-Benzenesulfonyl-4-piperazin-1-yl-1H-indole,
 2-benzenesulfonyl-4-(4-methyl-piperazin-1-yl)-1H-indole,
 2-(2,3-dichloro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
 2-(2,3-dichloro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole,
 15 2-benzenesulfonyl-1-methyl-4-piperazin-1-yl-1H-indole,
 2-(2-fluoro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
 2-(toluene-2-sulfonyl)-4-Piperazin-1-yl-1H-indole,
 2-(2-trifluoromethyl-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
 2-benzenesulfonyl-4-(3,5-dimethyl-piperazin-1-yl)-1H-indole,
 20 2-(3-bromo-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
 2-(thiophene-2-sulfonyl)-4-piperazin-1-yl-1H-indole,
 2-(2-fluoro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole,
 2-(2-fluoro-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole,
 2-(3-bromo-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole,
 25 2-benzenesulfonyl-4-piperidin-4-yl-1H-indole.

(xi) A process for producing a 2-substituted indole of the formula:



I

- 4 -

wherein

n is 0, 1 or 2;


p is 1 or 2;

R¹ is optionally substituted aryl or optionally substituted heteroaryl;

5 R² is a optionally substituted heterocyclyl;

R³ is hydrogen, alkyl, or $-C(=O)-R^5$, where R⁵ is alkyl, alkoxy, aryl, or aryloxy; and each R⁴ is independently hydrogen, hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxy carbonyl, alkyl carbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy;

said process comprising contacting a substituted indole of the formula:



II


15 wherein R^{2'} is a optionally substituted heterocyclyl, optionally protected with a protecting group; R^{3'} is alkyl or $-C(=O)-R^5$; each R^{4'} is independently hydrogen, hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxy carbonyl, alkyl carbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy,

20 optionally protected with a protecting group,

(a) with a base to produce a deprotonated indole; and

(b) contacting the deprotonated indole with a sulfonylating agent of the formula: $Y-SO_2-R^1$, where Y is halide, or a disulfide agent of the formula: $R^1-S-S-R^1$

25 to produce 2-substituted indole of the formula:



III

(c) optionally oxidizing the sulfur with an oxidizing agent; and

- 5 -

- (c) optionally oxidizing the sulfur with an oxidizing agent; and
 - (d) optionally removing the protecting group to produce the 2-substituted indole of formula I.
- (xii) The process of (xi), wherein Y is fluorine.
- 5 (xiii) A pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I of (i) to (x); and a pharmaceutically acceptable carrier.
- (xiv) Use of one or more compounds of any (i) to (x) for the manufacture of a medicament for the treatment or prevention of a disease state that is alleviated by 5-HT₆ agonists.
- 10 (xv) The use of (xiv), wherein the disease state comprises disorders of the CNS.
- (xvi) The use of (xv), wherein the disease state comprises psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic lateral sclerosis, Alzheimer's disease and Huntington's disease.
- 15 (xvii) The use of (xiv) wherein the disease state comprises disorder of the gastrointestinal.
- (xviii) The use of (xiv) wherein the disease state comprises obesity.

The present invention also provides methods for preparing, compositions comprising, and methods for using Compounds of Formula I.

- 20 Unless otherwise stated, the following terms used in this Application, including the specification and claims, have the definitions given below. It must be noted that, as used in the specification and the appended claims, the singular forms "a", "an," and "the" include plural referents unless the context clearly dictates otherwise.

"Agonist" refers to a compound that enhances the activity of another compound or receptor site.

- 25 "Alkyl" means the monovalent linear or branched saturated hydrocarbon moiety, consisting solely of carbon and hydrogen atoms, having from one to twelve, preferably one to four, carbon atoms. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl, sec-butyl, tert-butyl, pentyl, n-hexyl, octyl, dodecyl, and the like.

“Alkoxy” refers to a moiety of the formula -OR^a where R^a is alkyl as defined herein.

“Antagonist” refers to a compound that diminishes or prevents the action of another compound or receptor site.

“Aryl” means a monovalent cyclic aromatic hydrocarbon moiety consisting of a mono- or bicyclic aromatic ring. The aryl group can optionally be substituted with one, two or three, preferably one or two, substituents, wherein each substituent is independently hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy, unless otherwise specifically indicated. Examples of aryl moieties include, but are not limited to, optionally substituted phenyl and optionally substituted naphthyl, and the like.

“Aryloxy” refers to a moiety of the formula -OR^b where R^b is aryl as defined herein.

“Cycloalkyl” means a monovalent saturated carbocyclic moiety consisting of mono- or bicyclic rings. Cycloalkyl can optionally be substituted with one or more substituents, wherein each substituent is independently hydroxy, alkyl, alkoxy, halo, haloalkyl, amino, monoalkylamino, or dialkylamino, unless otherwise specifically indicated. Examples of cycloalkyl moieties include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.

“Disease state” means any disease, condition, symptom, or indication.

The terms “halo,” “halide,” and “halogen” are used interchangeably herein and refer to a substituent fluoro, chloro, bromo, or iodo, preferably fluoro or bromo.

“Haloalkyl” means alkyl as defined herein in which one or more hydrogen has been replaced with same or different halogen. Exemplary haloalkyls include -CH₂Cl, -CH₂CF₃, -CH₂CCl₃, perfluoroalkyl (e.g., -CF₃), and the like.

“Heteroaryl” means a monovalent mono-, bi-, or tri-cyclic aromatic moiety of 5 to 12 ring atoms containing one, two, three, or four ring heteroatoms selected from N, O, or S, the remaining ring atoms being C. The heteroaryl ring is optionally substituted independently with one or more substituents, preferably one or two substituents, selected from hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino,

alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy. More specifically the term heteroaryl includes, but is not limited to, pyridyl, furanyl, thiophenyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrazolyl, pyrimidinyl, benzofuranyl, isobenzofuranyl, benzothiazolyl, benzoisothiazolyl, 5 benzotriazolyl, indolyl, isoindolyl, benzoxazolyl, quinolyl, isoquinolyl, benzimidazolyl, benzisoxazolyl, benzothiophenyl, dibenzofuran, and benzodiazepin-2-one-5-yl, and the like.

“Heterocyclyl” means a monovalent saturated moiety, consisting of one to three rings, incorporating one, two, or three heteroatoms (chosen from nitrogen, oxygen or 10 sulfur). The heterocyclyl ring is optionally substituted independently with one or more substituents, preferably one or two substituents, selected from hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, 15 alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy. More specifically the term heterocyclyl includes, but is not limited to Examples of heterocyclyl moieties include, but are not limited to, morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, tetrahydropyranyl, azetidyl, and the like.

“Leaving group” means the group with the meaning conventionally associated with 20 it in synthetic organic chemistry, i.e., an atom or group displaceable under substitution reaction conditions. It should be appreciated that a particular leaving group depends on the reaction conditions including the atom to which the leaving group is attached to. For example, leaving groups for sulfonyl compounds include, but are not limited to, halogen, sulfonates, and the like.

25 “Modulator” means a molecule that interacts with a target. The interactions include, but are not limited to, agonist, antagonist, and the like, as defined herein.

“Optional” or “optionally” means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

30 “Inert organic solvent” or “inert solvent” means the solvent is inert under the conditions of the reaction being described in conjunction therewith, including for example, benzene, toluene, acetonitrile, tetrahydrofuran, N,N-dimethylformamide, chloroform, methylene chloride or dichloromethane, dichloroethane, diethyl ether, ethyl acetate, acetone, methyl ethyl ketone, methanol, ethanol, propanol, isopropanol, *tert*-

butanol, dioxane, pyridine, and the like. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert solvents.

“Pharmaceutically acceptable” means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary as well as human pharmaceutical use.

“Pharmaceutically acceptable salts” of a compound means salts that are pharmaceutically acceptable, as defined herein, and that possess the desired pharmacological activity of the parent compound. Such salts include:

acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, benzenesulfonic acid, benzoic, camphorsulfonic acid, citric acid, ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, hydroxynaphthoic acid, 2-hydroxyethanesulfonic acid, lactic acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, muconic acid, 2-naphthalenesulfonic acid, propionic acid, salicylic acid, succinic acid, tartaric acid, p-toluenesulfonic acid, trifluoroacetic acid, trimethylacetic acid, and the like; or

salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic or inorganic base. Acceptable organic bases include diethanolamine, ethanolamine, N-methylglucamine, triethanolamine, tromethamine, and the like. Acceptable inorganic bases include aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate and sodium hydroxide.

The preferred pharmaceutically acceptable salts are the salts formed from acetic acid, hydrochloric acid, sulfuric acid, methanesulfonic acid, maleic acid, phosphoric acid, tartaric acid, citric acid, sodium, potassium, calcium, zinc, and magnesium.

It should be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same acid addition salt.

“Prodrug” or “pro-drug” means a pharmacologically inactive form of a compound which must be metabolized in vivo, e.g., by biological fluids or enzymes, by a subject after administration into a pharmacologically active form of the compound in order to produce the desired pharmacological effect. The prodrug can be metabolized before absorption, during absorption, after absorption, or at a specific site. Although

metabolism occurs for many compounds primarily in the liver, almost all other tissues and organs, especially the lung, are able to carry out varying degrees of metabolism. Prodrug forms of compounds may be utilized, for example, to improve bioavailability, improve subject acceptability such as by masking or reducing unpleasant characteristics
5 such as bitter taste or gastrointestinal irritability, alter solubility such as for intravenous use, provide for prolonged or sustained release or delivery, improve ease of formulation, or provide site-specific delivery of the compound. Reference to a compound herein includes prodrug forms of a compound.

“Protective group” or “protecting group” means the group which selectively blocks
10 one reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning conventionally associated with it in synthetic chemistry. Certain processes of this invention rely upon the protective groups to block reactive nitrogen and/or oxygen atoms present in the reactants. For example, the terms “amino-protecting group” and “nitrogen protecting
15 group” are used interchangeably herein and refer to those organic groups intended to protect the nitrogen atom against undesirable reactions during synthetic procedures. Exemplary nitrogen protecting groups include, but are not limited to, trifluoroacetyl, acetamido, benzyl (Bn), benzyloxycarbonyl (carbobenzyloxy, CBZ), p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, *tert*-butoxycarbonyl (BOC), and
20 the like. Similarly, the term “hydroxy protecting group” refers to those organic groups intended to protect the oxygen atom of a hydroxyl group against undesirable reactions during synthetic procedures. Exemplary hydroxy protecting groups include, but are not limited to benzyl, silyl groups, tetrahydropyranyl, esters, and the like. The artisan in the art will know how to chose a group for the ease of removal and for the ability to
25 withstand the following reactions.

“Solvates” means solvent additions forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate, when the solvent is alcohol, the
30 solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one of the substances in which the water retains its molecular state as H₂O, such combination being able to form one or more hydrate.

“Subject” means mammals and non-mammals. Mammals means any member of the mammalia class including, but not limited to, humans; non-human primates such as
35 chimpanzees and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, and swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice, and guinea pigs; and the like. Examples of

- 10 -

non-mammals include, but are not limited to, birds, and the like. The term "subject" does not denote a particular age or sex.

"Therapeutically effective amount" means an amount of a compound that, when administered to a subject for treating a disease state, is sufficient to effect such treatment for the disease state. The "therapeutically effective amount" will vary depending on the compound, disease state being treated, the severity of the disease treated, the age and relative health of the subject, the route and form of administration, the judgment of the attending medical or veterinary practitioner, and other factors.

The terms "those defined above" and "those defined herein" when referring to a variable incorporates by reference the broad definition of the variable as well as preferred, more preferred and most preferred definitions, if any.

"Treating" or "treatment" of a disease state includes:

(i) preventing the disease state, i.e. causing the clinical symptoms of the disease state not to develop in a subject that may be exposed to or predisposed to the disease state, but does not yet experience or display symptoms of the disease state.

(ii) inhibiting the disease state, i.e., arresting the development of the disease state or its clinical symptoms, or


(iii) relieving the disease state, i.e., causing temporary or permanent regression of the disease state or its clinical symptoms.

The terms "treating", "contacting" and "reacting" when referring to a chemical reaction means adding or mixing two or more reagents under appropriate conditions to produce the indicated and/or the desired product. It should be appreciated that the reaction which produces the indicated and/or the desired product may not necessarily result directly from the combination of two reagents which were initially added, i.e., there may be one or more intermediates which are produced in the mixture which ultimately leads to the formation of the indicated and/or the desired product.

In general, the nomenclature used in this Application is based on AUTONOMTM v.4.0, a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature.

In one aspect, the present invention provides a compound of the formula:

- 11 -



I

a pharmaceutically acceptable salt or a prodrug thereof,

wherein

5 n is 0, 1 or 2; preferably n is 2;

p is 1 or 2, preferably p is 1;

R¹ is aryl or heteroaryl;

R² is heterocyclyl;

R³ is hydrogen, alkyl, or $-C(=O)-R^5$, where R⁵ is alkyl, alkoxy, aryl, or aryloxy; and

10 each R⁴ is independently hydrogen, hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxy carbonyl, alkyl carbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy; preferably R⁴ is hydrogen,
15 alkyl, alkoxy, halo, or haloalkyl.

It is to be understood that the scope of this invention encompasses not only the various isomers which may exist but also the various mixture of isomers which may be formed. Furthermore, the scope of the present invention also encompasses solvates and salts of Compounds of Formula I.

20 Preferably, R¹ is thienyl or phenyl which is optionally substituted with one or more substituents selected from the group consisting of alkyl, alkoxy, halo, haloalkyl, $-SO_2-R^a$, $-NR^aR^b$, $-NR^b-C(=O)-R^a$, where R^a is alkyl or aryl, and R^b is hydrogen or alkyl, and a mixture thereof. More preferably, R¹ is thien-2-yl, phenyl, 2,3-dichlorophenyl, 2-fluorophenyl, 2-methylphenyl, 2-trifluoromethylphenyl, or 3-bromophenyl.

25 Preferably, R² is optionally substituted piperazinyl or piperidinyl. More preferably, R² is optionally substituted piperazin-1-yl or piperidin-4-yl. Still more preferably, R² is piperazin-1-yl, 4-methylpiperazin-1-yl, 3,5-dimethylpiperazin-1-yl, N-methyl piperidin-4-yl or piperidin-4-yl. In one particular embodiment, R² is 4-methylpiperazin-1-yl.

- 12 -





Preferably, R³ is hydrogen or methyl.








Preferably, R⁴ is hydrogen.





Still further, combinations of the preferred groups described herein will form other preferred embodiments. For example, in one particularly preferred embodiment R¹ is phenyl, n is 2, R² is piperazin-1-yl, R³ is hydrogen or methyl, and R⁴ is hydrogen. In this manner, a variety of preferred compounds are embodied within the present invention.

Some of the representative Compounds of Formula I are shown in Table 1 below:

Table 1. Representative Compounds of Formula I:

#	Structure	Name	M+H
1		2-Benzenesulfonyl-4-piperazin-1-yl-1H-indole	342
2		2-Benzenesulfonyl-4-(4-methyl-piperazin-1-yl)-1H-indole	356
3		2-(2,3-Dichloro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole	410
4		2-(2,3-Dichloro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole	424

#	Structure	Name	M+H
5		2-Benzenesulfonyl-1-methyl-4-piperazin-1-yl-1H-indole	356
6		2-(2-Fluoro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole	360
7		2-(toluene-2-sulfonyl)-4-Piperazin-1-yl-1H-indole	356
8		2-(2-Trifluoromethyl-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,	410
9		2-Benzenesulfonyl-4-(3,5-dimethyl-piperazin-1-yl)-1H-indole	370
10		2-(3-Bromo-benzenesulfonyl)-4-piperazin-1-yl-1H-indole	422
11		2-(Thiophene-2-sulfonyl)-4-piperazin-1-yl-1H-indole	348

#	Structure	Name	M+H
12		2-(2-Fluoro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole	374
13		2-(2-Fluoro-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole	374
14		2-(3-Bromo-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole	434
15		2-Benzenesulfonyl-4-piperidin-4-yl-1H-indole	340

Another aspect of the present invention provides a composition comprising a therapeutically effective amount of a Compound of Formula I and a pharmaceutically acceptable carrier.

- 5 Yet another aspect of the present invention provides a method for treating a CNS disease state in a subject comprising administering to the subject a therapeutically effective amount of a Compound of Formula I. Preferably, the disease state comprises psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic lateral sclerosis, Alzheimer's
- 10 disease and Huntington's disease.

Still another aspect of the present invention provides a method for treating a disorder of the gastrointestinal tract in a subject comprising administering to the subject a therapeutically effective amount of a Compound of Formula I.

- 15 -

In addition another aspect of the present invention provides a method for treating obesity in subject comprising administering to the subject a therapeutically effective amount of a compound of Formula I.

Another aspect of the present invention provides a method for producing a
5 Compound of Formul


Compounds of the present invention can be made by a variety of methods depicted in the illustrative synthetic reaction schemes shown and described below.

The starting materials and reagents used in preparing these compounds generally are either available from commercial suppliers, such as Aldrich Chemical Co., or are
10 prepared by methods known to those skilled in the art following procedures set forth in references such as *Fieser and Fieser's Reagents for Organic Synthesis*; Wiley & Sons: New York, 1991, Volumes 1-15; *Rodd's Chemistry of Carbon Compounds*, Elsevier Science Publishers, 1989, Volumes 1-5 and Supplementals; and *Organic Reactions*, Wiley & Sons: New York, 1991, Volumes 1-40. The following synthetic reaction schemes are merely
15 illustrative of some methods by which the compounds of the present invention can be synthesized, and various modifications to these synthetic reaction schemes can be made and will be suggested to one skilled in the art having referred to the disclosure contained in this Application.

The starting materials and the intermediates of the synthetic reaction schemes can
20 be isolated and purified if desired using conventional techniques, including but not limited to, filtration, distillation, crystallization, chromatography, and the like. Such materials can be characterized using conventional means, including physical constants and spectral data.

Unless specified to the contrary, the reactions described herein preferably are
25 conducted under an inert atmosphere at atmospheric pressure at a reaction temperature range of from about -78 °C to about 150 °C, more preferably from about 0 °C to about 125 °C, and most preferably and conveniently at about room (or ambient) temperature, e.g., about 20 °C.

In one embodiment, Compounds of Formula I, are prepared by deprotonating a
30 substituted indole of the formula:



- 16 -


II

with a base and adding a sulfonylating agent of the formula: $Y-SO_2-R^1$, where R^1 , R^2 , and R^4 are those defined herein, $R^{3'}$ is alkyl or $-C(=O)-R^5$, where R^5 is that defined herein, and Y is a leaving group, preferably halide, and more preferably fluoride. It should be appreciated that when R^2 and/or R^4 has one or more acidic protons (relative to the base used), it should be protected with an appropriate protecting group. Suitable protecting groups for such acidic protons are well known to one skilled in the art and depends on the nature of the acidic proton, e.g., whether it is an amino proton or hydroxy proton, etc.

The base should be strong enough to deprotonate primarily the 2-position of the indole ring system. Such bases are well known to one skilled in the art and include organometallic compounds such as organolithiums, for example, tert-butyllithium, and Grignard reagents, for example, tert-butylmagnesium halide. Generally, the deprotonation reaction is conducted at a temperature of about 0 °C or below, preferably about -40 °C or below, and more preferably about -70 °C or below. Typically, the deprotonation reaction is conducted at about -75 °C.

Suitable sulfonylating agent include arylsulfonylhalides, for example, arylsulfonylfluorides. Arylsulfonylfluorides can be readily prepared from the corresponding arylsulfonylchlorides by treatment with a fluoride source, such as potassium fluoride or other suitable metallic fluoride compounds. Conversion of the arylsulfonylchloride to its corresponding fluoride derivative typically involves reacting the arylsulfonylchloride with potassium fluoride in an inert organic solvent, such as 1,4-dioxane. The reaction is generally carried out under refluxing conditions for a period of about 1 to about 48 hours, typically about 24 hours. Generally, excess potassium fluoride is used in the reaction, which can be readily removed during a work-up process by washing with water. The resulting arylsulfonylfluoride is typically used without any further purification.

Alternatively, Compounds of Formula I can also be prepared by reacting the deprotonated indole group with a disulfide reagent of the formula: $R^1-S-S-R^1$ to produce a thioether of the formula:



III

where R^1 , R^2 , $R^{3'}$, and R^4 are those defined herein. Typically, the disulfide reagent is added to the deprotonated indole at the same temperature in which the base is added. The reaction mixture is then stirred at that temperature for few minutes to few hours, typically from about 1 to 2 hours, and is allowed to gradually warm to room temperature.

5 The thioether compound of Formula III can be oxidized using an oxidizing agent to produce a corresponding sulfoxide and/or sulfone. Suitable oxidizing agents include *meta*-chloroperbenzoic acid (MCPBA), periodates, Oxone[®], as well as other sulfur oxidizing agents known to one skilled in the art. For example, the thioether III can be reacted with MCPBA by combining the two reagents at about 0 °C in an inert solvent,
10 such as dichloromethane, and stirring the mixture at room temperature for few hours. The excess MCPBA is typically removed by washing with an aqueous, preferably a basic aqueous, solution. Any undesired oxidation of nitrogen atom can be reduced by quenching the crude product with a phosphine compound, such as triphenylphosphine.

When R^1 or R^2 group contains a protecting group, or when $R^{3'}$ is a protecting
15 group, such protecting group can be removed after the synthesis using reaction conditions conventionally known to one skilled in the art. See, for example, *Protective Groups in Organic Synthesis*, 3rd edition, T.W. Greene and P.G.M. Wuts, John Wiley & Sons, New York, 1999, which is incorporated herein by reference in its entirety.

More specific details for producing Compounds of Formula I are described in the
20 Examples section.

Surprisingly and unexpectedly, the present inventors have found that indole compounds having a substituent $-S(O)_nR^1$ on the 2-position have a wide range of pharmacological activities that are not present in the corresponding indoles that do not have a such substituent on the 2-position. For example, the compounds of the invention
25 have selective 5-HT₆ receptor affinity and are useful in the treatment of certain CNS disorders, such as Parkinson's disease, Huntington's disease, anxiety, depression, manic depression, psychosis, epilepsy, obsessive compulsive disorders, migraine, Alzheimer's disease (enhancement of cognitive memory), sleep disorders, feeding disorders such as anorexia and bulimia, panic attacks, attention deficit hyperactivity disorder (ADHD),
30 attention deficit disorder (ADD), withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. In addition, compounds of the present invention are useful in the treatment of certain GI (gastrointestinal) disorders, such functional bowel disorder.

- 18 -

The pharmacology of the compounds of this invention was determined by art recognized procedures. The in vitro techniques for determining the affinities of test compounds at the 5-HT₆ receptor in radioligand binding and functional assays are described in Examples 4-6.

5 The present invention includes pharmaceutical compositions comprising at least one compound of the present invention, or an individual isomer, racemic or non-racemic mixture of isomers or a pharmaceutically acceptable salt or solvate thereof, together with at least one pharmaceutically acceptable carrier, and optionally other therapeutic and/or prophylactic ingredients.

10 In general, the compounds of the present invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. Suitable dosage ranges are typically 1-500 mg daily, preferably 1-100 mg daily, and most preferably 1-30 mg daily, depending upon numerous factors such as the severity of the disease to be treated, the age and relative
15 health of the subject, the potency of the compound used, the route and form of administration, the indication towards which the administration is directed, and the preferences and experience of the medical practitioner involved. One of ordinary skill in the art of treating such diseases will be able, without undue experimentation and in reliance upon personal knowledge and the disclosure of this Application, to ascertain a
20 therapeutically effective amount of the compounds of the present invention for a given disease.

In general, compounds of the present invention will be administered as pharmaceutical formulations including those suitable for oral (including buccal and sub-lingual), rectal, nasal, topical, pulmonary, vaginal, or parenteral (including
25 intramuscular, intraarterial, intrathecal, subcutaneous and intravenous) administration or in a form suitable for administration by inhalation or insufflation. The preferred manner of administration is generally oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction.

A compound or compounds of the present invention, together with one or more
30 conventional adjuvants, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds or principles, and the unit dosage forms may contain any suitable effective amount of the active ingredient
35 commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids,

powders, sustained release formulations, or liquids such as solutions, suspensions, emulsions, elixirs, or filled capsules for oral use; or in the form of suppositories for rectal or vaginal administration; or in the form of sterile injectable solutions for parenteral use. Formulations containing about one (1) milligram of active ingredient or, more broadly, about 0.01 to about one hundred (100) milligrams, per tablet, are accordingly suitable representative unit dosage forms.

The compounds of the present invention may be formulated in a wide variety of oral administration dosage forms. The pharmaceutical compositions and dosage forms may comprise a compound or compounds of the present invention or pharmaceutically acceptable salts thereof as the active component. The pharmaceutically acceptable carriers may be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about one (1) to about seventy (70) percent of the active compound. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier, providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges may be as solid forms suitable for oral administration.

Other forms suitable for oral administration include liquid form preparations including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions, or solid form preparations which are intended to be converted shortly before use to liquid form preparations. Emulsions may be prepared in solutions, for example, in aqueous propylene glycol solutions or may contain emulsifying agents, for example, such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known

suspending agents. Solid form preparations include solutions, suspensions, and emulsions, and may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

5 The compounds of the present invention may be formulated for parenteral administration (e.g., by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, for
10 example solutions in aqueous polyethylene glycol. Examples of oily or nonaqueous carriers, diluents, solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils (e.g., olive oil), and injectable organic esters (e.g., ethyl oleate), and may contain formulatory agents such as preserving, wetting, emulsifying or suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder
15 form, obtained by aseptic isolation of sterile solid or by lyophilization from solution for constitution before use with a suitable vehicle, e.g., sterile, pyrogen-free water.

 The compounds of the present invention may be formulated for topical administration to the epidermis as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily
20 base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also containing one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. Formulations suitable for topical administration in the mouth include lozenges comprising active agents in a flavored base, usually sucrose and acacia or
25 tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

 The compounds of the present invention may be formulated for administration as suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa
30 butter is first melted and the active component is dispersed homogeneously, for example, by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

 The compounds of the present invention may be formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in
35 addition to the active ingredient such carriers as are known in the art to be appropriate.

The compounds of the present invention may be formulated for nasal administration. The solutions or suspensions are applied directly to the nasal cavity by conventional means, for example, with a dropper, pipette or spray. The formulations may be provided in a single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomizing spray pump.

The compounds of the present invention may be formulated for aerosol administration, particularly to the respiratory tract and including intranasal administration. The compound will generally have a small particle size for example of the order of five (5) microns or less. Such a particle size may be obtained by means known in the art, for example by micronization. The active ingredient is provided in a pressurized pack with a suitable propellant such as a chlorofluorocarbon (CFC), for example, dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, or carbon dioxide or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by a metered valve. Alternatively the active ingredients may be provided in a form of a dry powder, for example a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). The powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of e.g., gelatin or blister packs from which the powder may be administered by means of an inhaler.

When desired, formulations can be prepared with enteric coatings adapted for sustained or controlled release administration of the active ingredient. For example, the compounds of the present invention can be formulated in transdermal or subcutaneous drug delivery devices. These delivery systems are advantageous when sustained release of the compound is necessary and when patient compliance with a treatment regimen is crucial. Compounds in transdermal delivery systems are frequently attached to a skin-adhesive solid support. The compound of interest can also be combined with a penetration enhancer, e.g., Azone (1-dodecylazacycloheptan-2-one). Sustained release delivery systems are inserted subcutaneously into the subdermal layer by surgery or injection. The subdermal implants encapsulate the compound in a lipid soluble membrane, e.g., silicone rubber, or a biodegradable polymer, e.g., polylactic acid.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and

- 22 -

powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.


Other suitable pharmaceutical carriers and their formulations are described in *Remington: The Science and Practice of Pharmacy* 1995, edited by E. W. Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania. Representative pharmaceutical formulations containing a compound of the present invention are described in Examples 6-12.

EXAMPLES

The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

Example 1 :

This example illustrates a method for producing Compounds of Formula I using the synthetic scheme outlined below:




Step 1

To a solution of the indole 1-1 (1.38 g, 4.6 mmol) in THF (35 mL) at room temperature was added di-tert-butyl dicarbonate (1.0 g, 4.6 mmol), followed by DMAP (56 mg, 0.46 mmol). The reaction mixture was allowed to stir at room temperature under nitrogen for 1 hour at which time the reaction mixture was concentrated *in vacuo*. The residue was partitioned between ethyl acetate (50 mL) and a saturated aqueous sodium bicarbonate solution (50 mL). The organic layer was washed with a saturated sodium bicarbonate solution (2 x 25 mL), and then brine (25 mL). The organic layer was dried (MgSO₄), filtered, and concentrated to afford a clear oil. The oil was crystallized in

- 23 -

hexanes to afford 1.65 g (89%) of boc-piperazine boc-indole (1-2) as a white solid. $(M+H)^+ = 402.2$.

Step 2a



- 5 Potassium fluoride (99%) (12 g, 216 mmol) was added to a solution of 2-fluorobenzenesulfonyl chloride (10 g, 51 mmol) in 1,4 dioxane (35 mL). The reaction mixture was refluxed for 24 hrs. then cooled to room temperature and poured into ice water (200 mL). The ice water was extracted with chloroform (3 x 50 mL). The combined chloroform layers were dried (MgSO₄), filtered, and concentrated to afford 2-
- 10 fluorobenzenesulfonyl fluoride (2a) as a colorless oil 6.7 g (74%). $(M+H)^+ = 177$

Step 2

- To a -75 °C solution of the bis-boc-indole 2 (500 mg, 1.25 mmol) in THF (25 mL) was slowly added t-BuLi (1.47 mL, 2.5 mmol). The reaction mixture was stirred for 30 minutes at which time 2-fluorobenzenesulfonyl fluoride 2a (222 mg, 1.5 mmol) was
- 15 added. The reaction was stirred for 1.5 h at which time the ice bath was removed and the reaction was allowed to warm over 45 min. The reaction mixture was quenched with a saturated solution of ammonium chloride (55 mL). The mixture was then extracted with ethyl acetate (3 x 20 mL). The combined ethyl acetate layers were washed with brine (25 mL). The organic layer was dried (MgSO₄), filtered, and concentrated to afford a yellow
- 20 oil. The oil was chromatographed over silica eluting with hexane:ethyl acetate (17:3) to afford 300 mg (43%) of 2-phenylsulfonyl-4-piperazin-1-yl indole (1-3).

Step 3

- 2-Phenylsulfonyl-4-piperazin-1-yl indole (1-3) (300 mg, 0.54 mmol) was dissolved in 1 M ethanolic HCl solution (30 mL) and warmed to reflux. After 2.5 h the reaction
- 25 mixture was cooled to room temperature and ether (30 mL) was added. The grey precipitate was collected to afford 150 mg (65%) of the dihydrochloride salt of the product (6). $(M+H)^+ = 360$.

Similarly, compounds 1, 3, 7, 8, 9, 10 and 11 of Table 1 above were synthesized in the same manner as above:

- 24 -

2-benzenesulfonyl-4-piperazin-1-yl-1H-indole,

2-(2,3-dichloro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,

2-(toluene-2-sulfonyl)-4-piperazin-1-yl-1H-indole,

2-(2-trifluoromethyl-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,

5 2-benzenesulfonyl-4-(3,5-dimethyl-piperazin-1-yl)-1H-indole,


2-(3-bromo-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,

2-(thiophene-2-sulfonyl)-4-piperazin-1-yl-1H-indole.

Example 2

10 This example illustrates a method for alkylating piperazine moiety of Compounds of Formula I.

2-benzenesulfonyl-4-(4-methyl-piperazin-1-yl)-1H-indole



To a solution of the free base of Compound 1 in Table 1 (269 mg, 0.79 mmol) and formaldehyde (30%) (0.3 mL, 3.97 mmol) in THF (20 mL) under inert atmosphere was added sodium triacetoxyborohydride (250 mg, 1.2 mmol) in a single portion. The reaction was allowed to stir at room temperature for 24 hrs at which time it was concentrated *in vacuo*. The residue was partitioned between 1 M sodium hydroxide (25 mL) and ethyl acetate (25 mL). The aqueous layer was extracted with ethyl acetate (2 x 15 mL). The combined organic layers were dried (MgSO_4), filtered and concentrated. The remaining oil was chromatographed over silica gel eluting with dichloromethane/methanol (5:1) to afford the product as a clear oil (140 mg, 0.39 mmol). The TFA salt of the oil was made by dissolving the oil in dichloromethane (10 mL) and adding TFA (90 mg, 0.79 mmol). The dichloromethane was concentrated and the remaining oil was crystallized from ethanol to afford 110 mg (39%) of Compound 2 of Table 1. $\text{M}+\text{H}=356$.

15
20
25

- 25 -

Similarly, compounds 4 and 12 of Table 1 above were synthesized in the same manner as above:


2-(2,3-Dichloro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole,

2-(2-Fluoro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole.

5 Example 3

This example illustrates another method for preparing Compounds of Formula I.

2-benzenesulfonyl-1-methyl-4-piperazin-1-yl-1H-indole



Step 1

10 To a solution of indole 1-1 (500 mg, 1.7 mmol) in THF (20 mL) under nitrogen was added potassium tert-butoxide (223 mg, 2.0 mmol). The reaction was allowed to stir for 30 min. at which time iodomethane (0.16 mL, 2.5 mmol) was added. The reaction was stirred for 3 hrs then quenched with water (20 mL) and extracted with ethyl acetate (2x30 mL). The ethyl acetate layer was washed with brine (25 mL), dried (MgSO₄),
 15 filtered, and concentrated to afford a green oil (540 mg, 1.6 mmol). The oil was chromatographed over silica eluting with ethyl acetate/hexanes (1:6) to afford 420 mg (80%) of compound 3-1 as a white solid. M+H=316.

Step 2

20 To a -75 °C solution of the N-methylindole compound 3-1 (330 mg, 1.05 mmol) in THF (25 mL) was slowly added t-BuLi (1.2 mL, 2.1 mmol). The reaction mixture was stirred for 30 minutes at -75 °C then warmed in an ice bath for 15 min. The reaction was re-cooled to -75 °C and benzenesulfonyl fluoride was added (0.2 mL, 1.6 mmol). The reaction was stirred cold for 1.5 h at which time the ice bath was removed and the reaction was allowed to warm to room temperature (45 min.). The reaction was
 25 quenched with a saturated solution of ammonium chloride (55 mL) and extracted with ethyl acetate (3 x 20 mL). The combined ethyl acetate layers were washed with brine

- 26 -

(25 mL), dried (MgSO₄), filtered and concentrated to afford a yellow oil. The oil was chromatographed over silica eluting with hexanes/ethyl acetate (4:1) to afford 190 mg (40%) of the product (3-2).

Step 3

- 5 The indole 3-2 (190 mg, 0.54 mmol) was dissolved in 1 M ethanolic HCl (30 mL) and warmed to reflux. After 2.5 hrs, the reaction mixture was cooled to room temperature and ether (30 mL) was added. The white precipitate was collected to afford 130 mg (87%) of the dihydrochloride salt of compound 5 of Table 1. M+H=356.

- Similarly, compounds 4 and 12 of Table 1 above were synthesized in the same
10 manner as above:

2-(2-Fluoro-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole,


2-(3-bromo-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole.

Example 4

- 15 This example illustrates a method for producing Compounds of Formula I wherein R² is piperidin-4-yl, using the synthetic scheme outlined below:

2-Benzenesulfonyl-4-piperidin-4-yl)-1H-indole

- 27 -

*Step1*

Di-tert-butyl dicarbonate (17g, 76.5 mmol) was added to a solution of 4-bromoindole 4-1 (10g, 51 mmol) and DMAP (0.62g, 0.51 mmol) in THF (120 mL). The reaction mixture was stirred for 15 minutes, and then stripped. The remaining oil was partitioned between ether and water. The ether layer was washed with 3 x 50 mL of saturated sodium bicarbonate and 1x 75 mL of brine. The ether layer was dried over magnesium sulfate, filtered and stripped. The remaining brown oil was chromatographed over silica gel eluting with 1:100 ethyl acetate: hexanes to afford 15 g (100% yield) of 4-bromo-indole-1-carboxylic acid-tert-butyl ester 4-2.

Step2

A solution of 4-bromo-indole-1-carboxylic acid-tert-butyl ester 4-2 (4.9g, 16.5 mmol) in THF (150 mL) under Argon was cooled to -70°C , and $n\text{-BuLi}$ (12.4 mL, 24.8 mmol) was added over 20 min. The reaction mixture was warmed to -5°C in an ice bath and was stirred at this temperature for 30 min. The mixture was cooled to -70°C and a solution of N-Boc-piperidone 4-3 (3.3 g, 16.5 mmol) in THF (10 mL) was added over 15

- 28 -

min. The reaction was stirred for 45 min at -70°C and was then warmed to room temperature. The reaction was quenched with a saturated solution of ammonium chloride (75 mL) and partitioned between water (25 mL) and ethyl acetate (75 mL). The organic layer was washed with water (50 mL) and brine (50 mL), then dried (MgSO_4),
5 filtered and concentrated. The remaining brown oil was chromatographed, eluting with ethyl acetate:hexanes (1:9) to afford 1.32 g (19 % yield) of 4-(1-*tert*-butoxycarbonyl-4-hydroxy-piperidin-4-yl)-indole-1-carboxylic acid *tert*-butyl ester 4-4.

Step 3

4-(1-*tert*-Butoxycarbonyl-4-hydroxy-piperidin-4-yl)-indole-1-carboxylic acid *tert*-
10 butyl ester 4-4 (1.32g, 3.17 mmol) was heated to 180°C under house vacuum until the bubbling ceased. The reaction mixture was cooled and solidified to afford 4-Hydroxy-4-(1-*H*-indol-4-yl)-piperidine-1-carboxylic acid *tert*-butyl ester (1.0g, 100% yield) 4-5.

Step 4

4-Hydroxy-4-(1-*H*-indol-4-yl)-piperidine-1-carboxylic acid *tert*-butyl ester 4-5
15 (1.0g, 3.17 mmol) was combined with pyridine (20 mL) and phosphorus oxychloride (0.7 mL, 7.4 mmol) at room temperature under nitrogen, and was stirred overnight. The reaction was partitioned between ethyl acetate (55mL) and water (55mL). The ethyl acetate layer was washed with water (2 X 30 mL) and brine (55 mL), then dried (MgSO_4), filtered and concentrated to afford (670 mg, 71%) of 4-(1*H* -Indol-4-yl)-3,6-dihydro-
20 2*H*-pyridine-1-carboxylic acid *tert*-butyl ester 4-6.

Step 5

A mixture of 4-(1*H* -Indol-4-yl)-3,6-dihydro-2*H*-pyridine-1-carboxylic acid *tert*-
butyl ester 4-6 (670 mg, 2.24 mmol) and 10%Pd/C (70 mg) in ethanol (75 mL) was
placed under H_2 atmosphere on the parr shaker at 50 psi, and was shaken overnight. The
25 reaction mixture was filtered through a plug of celite capped with a glass filter. The filtrate was concentrated to afford (650 mg, 96%) of 4-(1*H*-Indol-4-yl)-piperidine-1-carboxylic acid *tert*-butyl ester 4-7.

Step 6

To a solution of 4-(1*H*-Indol-4-yl)-piperidine-1- carboxylic acid *tert*-butyl ester 4-7
30 (650 mg, 2.16 mmol) in THF (75 mL) at room temperature di-*tert*-butyl dicarbonate (519 mg, 2.38 mmol) was added, followed by a catalytic amount of DMAP (5.6 mg, 0.046 mmol). The reaction mixture was allowed to stir at room temperature under nitrogen for 24 hours at which time the reaction mixture was stripped. The remaining residue was partitioned between ethyl acetate (50 mL) and saturated sodium bicarbonate (50mL). The
35 organic layer was washed with saturated sodium bicarbonate (2 x 25 mL), and with brine (25mL). The organic layer was dried (MgSO_4), filtered, and concentrated to afford a clear

- 29 -

oil. The oil was chromatographed eluting with hexanes/ethyl acetate (23:1) to afford 470 mg (54%) of 4-(1-*tert*-Butoxycarbonyl-piperidin-4-yl)-indole-1-carboxylic acid *tert*-butyl ester 4-8.

Step 7

5 To a -75 °C solution of 4-(1-*tert*-Butoxycarbonyl-piperidin-4-yl)-indole-1-carboxylic acid *tert*-butyl ester 4-8 (470 mg, 1.2 mmol) in THF (20 ml), was slowly added *t*-BuLi (1.6 mL, 2.25 mmol). The reaction mixture was stirred for 45 minutes at -75 °C, and the benzenesulfonyl fluoride 2a was added (0.22 mL, 1.8 mmol). The reaction was stirred for 2.5 h at which time the ice bath was removed and the reaction was warmed to
10 room temperature, then quenched with a saturated solution of ammonium chloride (25 mL). The mixture was then extracted with ethyl acetate (3 x 20 mL). The combined ethyl acetate layers were washed with brine (25mL), dried (MgSO₄), filtered and stripped to afford a yellow oil. The oil was chromatographed over silica eluting with hexanes/ethyl acetate (4:1) to afford (310 mg, 60%) of the product 2-Benzenesulfonyl-4-(1-*tert*-
15 butoxycarbonyl-piperidin-4-yl)-indole-1-carboxylic acid *tert*-butyl ester 4-9.

Step 8

2-Benzenesulfonyl-4-(1-*tert*-butoxycarbonyl-piperidin-4-yl)-indole-1-carboxylic acid *tert*-butyl ester 4-9 (310 mg, 0.57 mmol) was dissolved in 1M ethanolic HCl (30 mL) and warmed to reflux. After 2.5 h the reaction mixture was cooled to room temperature
20 and ether (30 mL) was added. The white precipitate was collected to afford 138 mg (64%) of the product 2-Benzenesulfonyl-4-piperidin-4-yl)-1*H*-indole 15.

Example 5

This example illustrates *in vitro* radioligand binding studies of Compound of
25 Formula I.

The binding activity of compounds of this invention *in vitro* was determined as follows. Duplicate determinations of ligand affinity are made by competing for binding of [³H]LSD in cell membranes derived from HEK293 cells stably expressing recombinant human 5-HT₆ receptor.

30 All determinations were made in assay buffer containing 50 mM Tris-HCl, 10 mM MgSO₄, 0.5 mM EDTA, 1 mM ascorbic acid, pH 7.4 at 37 °C, in a 250 microliter reaction volume. Assay tubes containing [³H] LSD (5 nM), competing ligand, and membrane were incubated in a shaking water bath for 60 min. at 37 °C, filtered onto Packard GF-B plates (pre-soaked with 0.3% PEI) using a Packard 96 well cell harvester and washed 3

- 30 -

times in ice cold 50 mM Tris-HCl. Bound [^3H] LSD was determined as radioactive counts per minute using Packard TopCount.

Displacement of [^3H]LSD from the binding sites was quantified by fitting concentration-binding data to a 4-parameter logistic equation:

$$\text{binding} = \text{basal} + \left(\frac{\text{Bmax} - \text{basal}}{1 + 10^{-\text{Hill}(\log[\text{ligand}] - \log \text{IC}_{50})}} \right)$$

where Hill is the Hill slope, [ligand] is the concentration of competing radioligand and IC_{50} is the concentration of radioligand producing half-maximal specific binding of radioligand. The specific binding window is the difference between the Bmax and the basal parameters.

- 10 Using the procedures of Example 4, compounds of Formula I were tested and found to be selective 5-HT₆ antagonists. Representative activities are shown in Table 2.

Table 2 Radioligand binding data

#	Name	pKi
1	2-Benzenesulfonyl-4-piperazin-1-yl-1H-indole	9.9
2	2-Benzenesulfonyl-4-(4-methyl-piperazin-1-yl)-1H-indole	9.9
5	2-Benzenesulfonyl-1-methyl-4-piperazin-1-yl-1H-indole	9.7
9	2-Benzenesulfonyl-4-(3,5-dimethyl-piperazin-1-yl)-1H-indole	9.1
12	2-(2-Fluoro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole	9.9
14	2-(3-Bromo-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole	9.3

Example 6

- 15 The cognition-enhancing properties of compounds of the invention may be in a model of animal cognition: the object recognition task model. Four months old male Wistar rats (Charles River, The Netherlands) were used. Compounds were prepared daily and dissolved in physiological saline and tested at three doses. Administration was always

- 31 -

given i.p. (injection volume 1 mL/kg) 60 minutes before T1. Scopolamine hydrobromide was injected 30 minutes after compound injection. Two equal testing groups were made of 24 rats and were tested by two experimenters. The testing order of doses was determined randomly. The experiments were performed using a double blind protocol.


- 5 All rats were treated once with each dose condition. The object recognition test was performed as described by Ennaceur, A., Delacour, J., 1988, A new one-trial test for neurobiological studies of memory in rats. 1: Behavioral data. *Behav. Brain Res.* 31, 47-59.

- While the present invention has been described with reference to the specific
10 embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective spirit and scope of the present invention. All such modifications
15 are intended to be within the scope of the claims appended hereto.

- 32 -

Claims

1. A compound of the formula:



I

- 5 or a pharmaceutically acceptable salt thereof,
wherein

n is 0, 1 or 2;

p is 1 or 2;

R¹ is optionally substituted aryl or optionally substituted heteroaryl;

10 R² is a optionally substituted heterocyclyl;

R³ is hydrogen, alkyl, or -C(=O)-R⁵, where R⁵ is alkyl, alkoxy, aryl, or
aryloxy; and

each R⁴ is independently hydrogen, hydroxy, cyano, alkyl, alkoxy,
thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl,
15 alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino,
alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl,
alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl,
alkylsulfonylamino or methylenedioxy.

2. The compound of claim 1,
20 wherein

n is 2;

R¹ is optionally substituted aryl or optionally substituted heteroaryl;

R² is a optionally substituted heterocyclyl;

R³ is hydrogen, alkyl;

25 and


R⁴ is hydrogen.

3. The compound according to Claim 2, wherein R² is optionally substituted
piperazin-1-yl or optionally substituted piperidin-4-yl.

- 33 -

4. The compound according to Claim 3, wherein R² is piperazin-1-yl, 4-methylpiperazin-1-yl, 3,5-dimethylpiperazin-1-yl, N-methyl piperidin-4-yl or piperidin-4-yl.
5. The compound according to Claim 4, wherein R² is piperazin-1-yl, 4-methylpiperazin-1-yl or piperidin-4-yl.
6. The compound according to Claim 2, wherein R¹ is optionally substituted phenyl or optionally substituted thienyl.
7. The compound according to Claim 6, wherein R¹ is thien-2-yl or phenyl which is optionally substituted with alkyl, halo, or haloalkyl.
- 10 8. The compound according to Claim 7, wherein R¹ is phenyl, 2,3-dichlorophenyl, 2-fluorophenyl, 2-methylphenyl, 2-trifluoromethylphenyl, 3-bromophenyl, or thien-2-yl.
9. The compound according to Claim 2, wherein R³ is hydrogen or methyl.
10. The compounds of Claim 2, which are:
 - 15 2-Benzenesulfonyl-4-piperazin-1-yl-1H-indole,
2-benzenesulfonyl-4-(4-methyl-piperazin-1-yl)-1H-indole,
2-(2,3-dichloro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
2-(2,3-dichloro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole,
2-benzenesulfonyl-1-methyl-4-piperazin-1-yl-1H-indole,
 - 20 2-(2-fluoro-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
2-(toluene-2-sulfonyl)-4-Piperazin-1-yl-1H-indole,
2-(2-trifluoromethyl-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
2-benzenesulfonyl-4-(3,5-dimethyl-piperazin-1-yl)-1H-indole,
2-(3-bromo-benzenesulfonyl)-4-piperazin-1-yl-1H-indole,
 - 25 2-(thiophene-2-sulfonyl)-4-piperazin-1-yl-1H-indole,
2-(2-fluoro-benzenesulfonyl)-4-(4-methyl-piperazin-1-yl)-1H-indole,
2-(2-fluoro-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole,
2-(3-bromo-benzenesulfonyl)-1-methyl-4-piperazin-1-yl-1H-indole,
2-benzenesulfonyl-4-piperidin-4-yl-1H-indole.
- 30 11. A process for producing a 2-substituted indole of the formula:

- 34 -



I

wherein

n is 0, 1 or 2;

5 p is 1 or 2;


R¹ is optionally substituted aryl or optionally substituted heteroaryl;

R² is a optionally substituted heterocyclyl;

R³ is hydrogen, alkyl, or -C(=O)-R⁵, where R⁵ is alkyl, alkoxy, aryl, or aryloxy; and

10 each R⁴ is independently hydrogen, hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl,
15 alkylsulfonylamino or methylenedioxy;

said process comprising contacting a substituted indole of the formula:




II

wherein R²' is a optionally substituted heterocyclyl, optionally protected with a protecting
20 group; R³' is alkyl or -C(=O)-R⁵; each R⁴' is independently hydrogen, hydroxy, cyano, alkyl, alkoxy, thioalkyl, alkylthio, halo, haloalkyl, hydroxyalkyl, nitro, alkoxycarbonyl, alkylcarbonyl, alkylsulfonyl, arylsulfonyl, haloalkylsulfonyl, amino, alkylamino, dialkylamino, alkyl(aryl)amino, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl(alkylamino), alkylaminosulfonyl, alkylsulfonylamino or methylenedioxy,
25 optionally protected with a protecting group,

(i) with a base to produce a deprotonated indole; and

- 35 -

(ii) contacting the deprotonated indole with a sulfonylating agent of the formula: $Y-SO_2-R^1$, where Y is halide, or a disulfide agent of the formula: $R^1-S-S-R^1$ to produce 2-substituted indole of the formula:



III

5

(iii) optionally oxidizing the sulfur with an oxidizing agent; and

(iv) optionally removing the protecting group to produce the 2-substituted indole of formula I.

12. The process of Claim 11, wherein Y is fluorine.

10 13. A pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I of Claim 1 to 10; and a pharmaceutically acceptable carrier.

14. Use of one or more compounds of any claim 1 to 10 for the manufacture of a medicament for the treatment or prevention of a disease state that is alleviated by 5-HT6 agonists.

15 15. The use of claim 14, wherein the disease state comprises disorders of the CNS.

16. The use of Claim 15, wherein the disease state comprises psychoses, schizophrenia, manic depressions, neurological disorders, memory disorders, attention deficit disorder, Parkinson's disease, amyotrophic lateral sclerosis, Alzheimer's disease
20 and Huntington's disease.

17. The use of claim 14 wherein the disease state comprises disorder of the gastrointestinal.

18. The use of claim 14 wherein the disease state comprises obesity.

19. The invention as hereinbefore described.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/09969

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/30 A61K31/404 A61P43/00 C07D401/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02 32863 A (BIOVITRUM AB) 25 April 2002 (2002-04-25) page 2, line 10 - line 16; claim 1 ----	1,13,14
A	WO 02 41889 A (SMITHKLINE BEECHAM PLC) 30 May 2002 (2002-05-30) page 1, line 19 - line 25; claim 1 ----	1,13,14
A	WO 02 36562 A (AMERICAN HOME PRODUCTS CORPORATION) 10 May 2002 (2002-05-10) page 1; claim 1 -----	1,13,14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

11 February 2004

Date of mailing of the international search report

23/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/09969

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0232863	A	25-04-2002	AU 9619301 A	29-04-2002
			BR 0114552 A	01-07-2003
			CA 2422717 A1	25-04-2002
			EP 1326830 A1	16-07-2003
			NO 20031589 A	28-05-2003
			WO 0232863 A1	25-04-2002
			US 2002165251 A1	07-11-2002
WO 0241889	A	30-05-2002	AU 2071502 A	03-06-2002
			WO 0241889 A2	30-05-2002
			EP 1335722 A2	20-08-2003
WO 0236562	A	10-05-2002	AU 2005102 A	15-05-2002
			BR 0115102 A	30-09-2003
			CA 2426031 A1	10-05-2002
			EP 1343756 A2	17-09-2003
			NO 20031977 A	30-06-2003
			US 2002115670 A1	22-08-2002
			WO 0236562 A2	10-05-2002